



TITLE:

AN EXPLICIT INTEGRAL
REPRESENTATION OF WHITTAKER
FUNCTIONS FOR THE
REPRESENTATIONS OF THE
DISCRETE SERIES. : THE CASE OF
 $Sp(2; \mathbb{R})$

AUTHOR(S):

ODA, TAKAYUKI

CITATION:

ODA, TAKAYUKI. AN EXPLICIT INTEGRAL REPRESENTATION OF WHITTAKER FUNCTIONS FOR THE REPRESENTATIONS OF THE DISCRETE SERIES. : THE CASE OF $Sp(2; \mathbb{R})$. 数理解析研究所講究録 1992, 792: 151-171

ISSUE DATE:

1992-06

URL:

<http://hdl.handle.net/2433/82701>

RIGHT:

AN EXPLICIT INTEGRAL REPRESENTATION
OF WHITTAKER FUNCTIONS
FOR THE REPRESENTATIONS OF THE DISCRETE SERIES.
- THE CASE OF $Sp(2; \mathbb{R})$ -

TAKAYUKI ODA

1-st draft (α test)

Introduction.

We shall prove an explicit integral formula for the Whittaker function associated to the highest weight vector in the representation space of the minimal K -type of a discrete series representation with the maximal Gelfand-Kirillov dimension for the real symplectic group $Sp(2; \mathbb{R})$ of rank 2.

Let us explain the basic idea of this paper. Consider the case $G = SL_2(\mathbb{R})$. Put $N = \left\{ \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \mid x \in \mathbb{R} \right\}$, and let $\eta : \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \mapsto \exp(2\pi i c x)$ ($c \in \mathbb{R}$) be a non-trivial unitary character of N . Let $C_\eta^\infty(N \backslash G)$ be the space of C^∞ -functions φ satisfying $\varphi(n g) = \eta(n) \varphi(g)$ ($\forall (n, g) \in N \times G$).

For an irreducible unitary representation (π, H_π) of G , we denote by H_π^∞ the space of smooth vectors in G . When (π, H_π) is a principal series representation of $SL_2(\mathbb{R})$, the image of a vector in H_π^∞ with respect to a unique continuous intertwining operator from H_π^∞ to $C_\eta^\infty(N \backslash G)$ is represented by the modified Bessel function, i.e. the Whittaker function, if it is restricted to the split torus $A = \left\{ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \mid a \in \mathbb{R}, a > 0 \right\}$.

However when (π, H_π) is a discrete series representation of formal degree $k - 1$ of $SL_2(\mathbb{R})$, then the image of minimal K -type vector of H_π with respect to the intertwining operator from H_π^∞ to $C_\eta^\infty(N \backslash G)$ (if it exists), is written by $\text{const. } a^k e^{-2\pi|c| \cdot a^2}$ on A (cf. Jacquet-Langlands [J-L]).

Thus as special functions on A , the functions realizing the Whittaker model of the discrete series representations of $SL_2(\mathbb{R})$ are “degenerate” elementary functions, much simpler than those of the principal series representations.

We hope similar phenomena occur in higher rank groups. The purpose of this paper is to confirm this for the case $G = Sp(2; \mathbb{R})$. Let us explain the contents of this paper.

In the first place, we shall compute explicitly the partial differential equation for the radial part of the above Whittaker function. We follow the method of Yamashita [Y-I] [Y-II] who discussed the case $G = SU(2, 2)$.

In §1, we recall basic notation for the structure of $Sp(2; \mathbb{R})$ and associated Lie algebras.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

§2 reviews the Harish-Chandra parametrization of the representations of discrete series for $Sp(2; \mathbb{R})$. In §3, we recall the representation of $U(2)$, and in §4 the characters of the maximal unipotent subgroup of $Sp(2; \mathbb{R})$. In §5 – §8, we write down explicitly the system of partial differential equations characterizing the Whittaker functions of the minimal K -type of a discrete series representation.

New parts different from [Y-I], [Y-II] are Proposition (8.1) and §9. §9 contains the main result of this paper: an explicit integral expression of the Whittaker function of the highest weight vector of the minimal K -type of a discrete series representation of $Sp(2; \mathbb{R})$.

The author thanks to Professors T. Oshima and N. Wallach for educational conversations on the representation theory of real reductive groups in various occasions, to Professor H. Matsumoto for communications on the theory of Whittaker models, and to Professor T. Miwa for an assist to solve partial differential equations.

INTEGRAL REPRESENTATION OF WHITTAKER FUNCTION

§1 Basic Notations, and the structure of Lie groups and algebras.

In this section, we determine basic notations on the symplectic group of degree 2, its maximal compact subgroup and associated Lie algebras.

⟨Lie groups⟩

Let $M_4(\mathbb{R})$ be the space of real 4×4 matrices. Put $J = \begin{pmatrix} 0 & 1_2 \\ -1_2 & 0 \end{pmatrix} \in M_4(\mathbb{R})$, where 1_2 is a unit matrix of size 2. The symplectic group $Sp(2; \mathbb{R})$ of degree 2 is given by

$$Sp(2; \mathbb{R}) = \{g \in M_4(\mathbb{R}) \mid {}^t g J g = J, \det(g) = 1\}.$$

Here ${}^t g$ denotes the transpose of the matrix g , and $\det(g)$ the determinant of g . A maximal compact group K of $G = Sp(2; \mathbb{R})$ is given by

$$K = \left\{ \begin{pmatrix} A & B \\ -B & A \end{pmatrix} \in Sp(2; \mathbb{R}) \mid A, B \in M_2(\mathbb{R}) \right\},$$

which is isomorphic to the unitary group

$$U(2) = \{g \in GL(2; \mathbb{C}) \mid {}^t \bar{g} \cdot g = 1_2\}$$

of size 2 via a homomorphism

$$\begin{pmatrix} A & B \\ -B & A \end{pmatrix} \in K \longmapsto A + \sqrt{-1}B \in U(2).$$

⟨Lie algebras⟩

The Lie algebra of G is given by

$$\mathfrak{g} = \mathfrak{sp}(2; \mathbb{R}) = \{X \in M_4(\mathbb{R}) \mid JX + {}^t X J = 0\},$$

and that of K is given by

$$\mathfrak{k} = \left\{ X = \begin{pmatrix} A & B \\ -B & A \end{pmatrix} \mid A, B \in M_2(\mathbb{R}); {}^t A = -A, {}^t B = B \right\}.$$

The Cartan involution for \mathfrak{k} is given by

$$\theta(X) = -{}^t X \quad \text{for } X \in \mathfrak{g}.$$

Hence the subspace

$$\mathfrak{p} = \{X \in \mathfrak{g} \mid \theta(X) = X\} = \left\{ \begin{pmatrix} A & B \\ B & -A \end{pmatrix} \mid {}^t A = A, {}^t B = B; A, B \in M_2(\mathbb{R}) \right\}$$

given a Cartan decomposition

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}.$$

The linear map

$$\begin{pmatrix} A & B \\ -B & A \end{pmatrix} \in \mathfrak{k} \longmapsto A + \sqrt{-1}B \in \mathfrak{u}(2)$$

defines an isomorphism of Lie algebras from k to the unitary Lie algebras

$$\mathfrak{u}(2) = \{C \in M_2(\mathbb{C}) \mid {}^t\bar{C} + C = 0\}$$

of degree 2.

An \mathbb{R} -basis of $\mathfrak{u}(2)$ is given by

$$\sqrt{-1} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \sqrt{-1} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, Y = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, Y' = \sqrt{-1} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

Let $\mathfrak{u}(2)_{\mathbb{C}} = \mathfrak{u}(2) \otimes_{\mathbb{R}} \mathbb{C}$ be the complexification of $\mathfrak{u}(2)$. Then a basis of $\mathfrak{u}(2)_{\mathbb{C}}$ is given by

$$Z = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad H' = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

$$X = \frac{1}{2}(Y - \sqrt{-1}Y') = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \bar{X} = \frac{1}{2}(-Y - \sqrt{-1}Y') = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

Then $\{H', X, \bar{X}\}$ is a \mathfrak{sl}_2 -triple, i.e.

$$[H', X] = 2X; \quad [H', \bar{X}] = -2\bar{X}; \quad [X, \bar{X}] = H'.$$

Via the isomorphism $\mathfrak{k}_{\mathbb{C}} \xrightarrow{\sim} \mathfrak{u}_{\mathbb{C}}$, the preimage of the above basis of $\mathfrak{u}_{\mathbb{C}}$ is given by

$$Z = (-\sqrt{-1}) \left(\begin{array}{c|c} & 1 \\ \hline -1 & 1 \\ \hline & -1 \end{array} \right); \quad H' = (-\sqrt{-1}) \left(\begin{array}{c|c} & 1 \\ \hline -1 & -1 \\ \hline & 1 \end{array} \right);$$

$$Y = \left(\begin{array}{c|c} 0 & 1 \\ \hline -1 & 0 \\ \hline & 0 & 1 \\ \hline & -1 & 0 \end{array} \right); \quad Y' = \left(\begin{array}{c|c} & 1 \\ \hline & 1 \\ \hline -1 & 1 \\ \hline & -1 \end{array} \right).$$

From now on we use the convention that unwritten components of a matrix are zero. Now we fix a compact Cartan subalgebra \mathfrak{h} of \mathfrak{g} by

$$\mathfrak{h} = \mathbb{R}(\sqrt{-1}Z) + \mathbb{R}(\sqrt{-1}H').$$

Write $T_+ = \sqrt{-1}Z$ and $T_- = \sqrt{-1}H'$, and set

$$T_1 = \frac{1}{2}(T_+ + T_-) \text{ and } T_2 = \frac{1}{2}(T_+ - T_-).$$

Put

$$H'_1 = \frac{1}{2}(Z + H'), \quad H'_2 = \frac{1}{2}(Z - H').$$

Then $T_1 = \sqrt{-1}H'_1$, $T_2 = \sqrt{-1}H'_2$, and

$$T_1 = \left(\begin{array}{c|c} & 1 \\ \hline -1 & 0 \\ \hline & 0 \end{array} \right), \quad T_2 = \left(\begin{array}{c|c} & 0 \\ \hline 0 & 1 \\ \hline & -1 \end{array} \right) \in \mathfrak{h}.$$

INTEGRAL REPRESENTATION OF WHITTAKER FUNCTION

(Root system)

We consider a root space decomposition of \mathfrak{g} with respect to \mathfrak{h} . For a linear form $\beta : \mathfrak{h} \rightarrow \mathbb{C}$, we write $\beta(T_i) = \beta_i \in \mathbb{C}$. For each $\beta \in \mathfrak{h}^* = \text{Hom}(\mathfrak{h}, \mathbb{C})$, set

$$\mathfrak{g}_\beta = \{X \in \mathfrak{g}_{\mathbb{C}} = \mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C} \mid [H, X] = \beta(H)X, \forall H \in \mathfrak{h}\}.$$

Then the roots of $(\mathfrak{g}, \mathfrak{h})$ is given by

$$\begin{aligned} \Sigma &= \{\beta = (\beta_1, \beta_2) \mid \mathfrak{g}_\beta \neq 0, \beta \neq 0\} \\ &= \sqrt{-1}\{\pm(2, 0), \pm(0, 2), \pm(1, 1), \pm(1, -1)\}. \end{aligned}$$

We determine a root vector X_β in \mathfrak{g}_β , i.e. a generator of \mathfrak{g}_β by the following table.

$-\sqrt{-1}\beta$	$(2, 0)$	$(1, 1)$	$(0, 2)$	$(1, -1)$
X_β	$\left(\begin{array}{c c} 1 & i \\ \hline 0 & 0 \\ \hline i & -1 \\ 0 & 0 \end{array}\right)$	$\left(\begin{array}{c c} 1 & i \\ \hline 1 & i \\ \hline i & -1 \end{array}\right)$	$\left(\begin{array}{c c} 0 & 0 \\ \hline 1 & i \\ \hline 0 & 0 \\ i & -1 \end{array}\right)$	$\left(\begin{array}{c c} 1 & -i \\ \hline -1 & i \\ \hline i & 1 \\ -i & -1 \end{array}\right)$
$X_{-\beta}$	$\left(\begin{array}{c c} 1 & -i \\ \hline 0 & 0 \\ \hline -i & -1 \\ 0 & 0 \end{array}\right)$	$\left(\begin{array}{c c} 1 & -i \\ \hline 1 & -i \\ \hline -i & -1 \end{array}\right)$	$\left(\begin{array}{c c} 0 & 0 \\ \hline 1 & -i \\ \hline 0 & 0 \\ -i & -1 \end{array}\right)$	$\left(\begin{array}{c c} 1 & i \\ \hline -1 & -i \\ \hline -i & 1 \\ i & -1 \end{array}\right)$

Then

$$\mathfrak{k}_{\mathbb{C}} = \mathfrak{h}_{\mathbb{C}} + \mathbb{C}X_{(1, -1)} + \mathbb{C}X_{(-1, 1)},$$

and set

$$\begin{aligned} \mathfrak{p}_+ &= \mathbb{C}X_{(2, 0)} + \mathbb{C}X_{(1, 1)} + \mathbb{C}X_{(0, 2)} \\ &= \left\{X = \begin{pmatrix} X_1 & iX_1 \\ iX_1 & -X_1 \end{pmatrix} \mid X_1 \in M_2(\mathbb{C})\right\}, \end{aligned}$$

and

$$\begin{aligned} \mathfrak{p}_- &= \mathbb{C}X_{-(2, 0)} + \mathbb{C}X_{-(1, 1)} + \mathbb{C}X_{-(0, 2)} \\ &= \left\{X = \begin{pmatrix} X_1 & -iX_1 \\ -iX_1 & -X_1 \end{pmatrix} \mid X_1 \in M_2(\mathbb{C})\right\}. \end{aligned}$$

Then

$$\mathfrak{g}_{\mathbb{C}} = \mathfrak{k}_{\mathbb{C}} \oplus \mathfrak{p}_+ \oplus \mathfrak{p}_-.$$

For each root $\beta = (\beta_1, \beta_2)$, we put

$$\|\beta\| = \sqrt{|\beta_1|^2 + |\beta_2|^2}.$$

Then $\|\beta\|^2 = 4$ or $= 2$.

Then set

$$\{c \cdot \|\beta\|(X_\beta + X_{-\beta}), c \cdot \sqrt{-1}\|\beta\|(X_\beta - X_{-\beta}) \quad (\beta \in \Sigma_n^+)\}$$

forms an orthonormal basis of $\mathfrak{p} = \mathfrak{p}_{\mathbb{R}}$ with respect to the Killing form for some points constant c . Here $\Sigma_n^+ = \{(2, 0), (1, 1), (0, 2)\}$ is the set of non-compact positive roots. $\Sigma_c^+ = \{(1, -1)\}$ is the set of compact positive roots. $\Sigma_c = \Sigma_c^+ \cup (-\Sigma_c^+)$ and $\Sigma_n = \Sigma_n^+ \cup (-\Sigma_n^+)$ are the set of compact roots and the set of non-compact roots, respectively.

(Iwasawa decomposition)

We choose a maximal abelian subalgebra \mathfrak{a} of \mathfrak{p} given by

$$\mathfrak{a} = \left\{ \left(\begin{array}{c|c} A & 0 \\ \hline 0 & -A \end{array} \right) \mid A = \text{diag}(t_1, t_2) \quad (t_1, t_2 \in \mathbb{R}) \right\}.$$

Here $\text{diag}(t_1, t_2)$ is a diagonal matrix with $(1, 1)$ -entry t_1 and $(2, 2)$ -entry t_2 . Set

$$H_1 = \left(\begin{array}{c|c} 1 & \\ \hline 0 & -1 \\ \hline & 0 \end{array} \right) \quad \text{and} \quad H_2 = \left(\begin{array}{c|c} 0 & \\ \hline 1 & 0 \\ \hline & -1 \end{array} \right).$$

Then $\{H_1, H_2\}$ forms a basis of \mathfrak{a} .

(Root system of $(\mathfrak{g}, \mathfrak{a})$)

Let $\{e_1 = (1, 0), e_2 = (0, 1)\}$ be a standard basis of the 2-dimensional Euclidean plane \mathbb{R}^2 . Then the root system Ψ of $(\mathfrak{g}, \mathfrak{a})$ is given by

$$\Psi = \{\pm 2e_1, \pm 2e_2, \pm e_1 \pm e_2\}$$

A positive root system Ψ_+ is fixed by

$$\Psi_+ = \{2e_1, 2e_2, e_1 + e_2, e_1 - e_2\}.$$

Put

$$\mathfrak{n} = \sum_{\alpha \in \Psi_+} \mathfrak{g}_\alpha.$$

Then it is a nilradical of a minimal parabolic subalgebra. We choose generators E_α of \mathfrak{g}_α ($\alpha \in \Psi_+$) as follows.

$$\begin{aligned} E_{2e_1} &= \left(\begin{array}{c|cc} & 1 & 0 \\ \hline & 0 & 0 \\ \hline & & \end{array} \right); & E_{e_1+e_2} &= \left(\begin{array}{c|cc} & 0 & 1 \\ \hline & 1 & 0 \\ \hline & & \end{array} \right); \\ E_{2e_2} &= \left(\begin{array}{c|cc} & 0 & 0 \\ \hline & 0 & 1 \\ \hline & & \end{array} \right); & E_{e_1-e_2} &= \left(\begin{array}{cc|cc} 0 & 1 & & \\ 0 & 0 & & \\ \hline & & 0 & 0 \\ & & -1 & 0 \end{array} \right). \end{aligned}$$

The Iwasawa decomposition associated to $(\mathfrak{a}, \mathfrak{n})$ is given by

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{a} \oplus \mathfrak{n}.$$

In $\mathfrak{g}_{\mathbb{C}}$, the Iwasawa decomposition of the root vectors $\{X_\beta; \beta \in \Sigma\}$ are given as follows.

INTEGRAL REPRESENTATION OF WHITTAKER FUNCTION

Lemma (1.1).

$$\begin{aligned}
X_{(2,0)} &= H'_1 + H_1 + 2\sqrt{-1}E_{2e_1}; & X_{(-2,0)} &= -H'_1 + H_1 - 2\sqrt{-1}E_{2e_1}; \\
X_{(1,1)} &= 2 \cdot \overline{X} + 2 \cdot E_{e_1-e_2} + 2\sqrt{-1}E_{e_1+e_2}; \\
X_{(-1,-1)} &= -2 \cdot X + 2 \cdot E_{e_1-e_2} - 2\sqrt{-1}E_{e_1+e_2}; \\
X_{(0,2)} &= H'_2 + H_2 + 2\sqrt{-1}E_{2e_2}; & X_{(0,-2)} &= -H'_2 + H_2 - 2\sqrt{-1}E_{2e_2}.
\end{aligned}$$

■

Proof) A direct computation.

§2 Parametrization of the representation of the discrete series.

Consider a compact Cartan subgroup of G

$$\exp(\mathfrak{h}) = \left\{ \left(\begin{array}{cc|cc} \cos \theta_1 & & \sin \theta_1 & \\ & \cos \theta_2 & & \sin \theta_2 \\ \hline -\sin \theta_1 & & \cos \theta_1 & \\ & -\sin \theta_2 & & \cos \theta_2 \end{array} \right) \mid \theta_1, \theta_2 \in \mathbb{R} \right\}$$

corresponding to \mathfrak{h} . Then the characters are given by

$$\left(\begin{array}{cc|cc} \cos \theta_1 & & \sin \theta_1 & \\ & \cos \theta_2 & & \sin \theta_2 \\ \hline -\sin \theta_1 & & \cos \theta_1 & \\ & -\sin \theta_2 & & \cos \theta_2 \end{array} \right) \mapsto \exp\{\sqrt{-1}(m_1\theta_1 + m_2\theta_2)\} \in \mathbb{C}^*.$$

Here m_1, m_2 are some integers. The derivation of these characters determines an integral structure of $\mathfrak{h}^* = \text{Hom}(\mathfrak{h}, \mathbb{C})$, the weight lattice.

The set of compact positive roots is given by $\Sigma_c^+ = \{(1, -1)\}$. Hence the set of dominant weight are given by $\{(\lambda_1, \lambda_2) \in \mathbb{Z}^{\oplus 2} \mid \lambda_1 \geq \lambda_2\}$. In order to parametrize the representation of the discrete series of $\text{Sp}(2; \mathbb{R})$, we first enumerate all the positive root systems compatible to Σ_c^+ . There are four such positive root systems:

- (I) : $\Sigma_I^+ = \{(1, -1), (2, 0), (1, 1), (0, 2)\};$
- (II) : $\Sigma_{II}^+ = \{(1, -1), (1, 1), (2, 0), (0, -2)\};$
- (III) : $\Sigma_{III}^+ = \{(1, -1), (2, 0), (0, -2), (-1, -1)\};$
- (IV) : $\Sigma_{IV}^+ = \{(1, -1), (-2, 0), (-1, -1), (0, -2)\}.$

Let J be a variable running over the set of indices $\{I, II, III, IV\}$. Then we write $\Sigma_{J,n}^+ = \Sigma_J^+ - \Sigma_c^+$ for the set of non-compact positive roots for each index J .

Define a subset Ξ_J of dominant weights by

$$\Xi_J = \{\Lambda = (\Lambda_1, \Lambda_2) \text{ dominant w.r.t. } \Sigma_c^+ \mid \langle \Lambda, \beta \rangle > 0, \forall \beta \in \Sigma_{J,n}^+\}.$$

Then the set $\bigcup_{J=I}^{IV} \Xi_J$ gives the Harish-Chandra parametrization of the representation of the discrete series for $\text{Sp}(2; \mathbb{R})$. Let π_Λ be the associated representation of G for $\Lambda \in \bigcup_{J=I}^{IV} \Xi_J$. The K -types of $\pi_\Lambda|_K$ is described by the formula of Blattner proved finally by Hecht-Schmid []. Among others the minimal K -type of π_Λ is given by $\lambda_{\min} = \Lambda - \rho_c + \rho_n$. Here ρ_c and ρ_n are the half of the sum of compact positive roots and non-compact positive roots. Here is a table of λ_{\min} .

type J	I,	II,	III,	IV
λ_{\min}	$(\Lambda_1 + 1, \Lambda_2 + 2),$	$(\Lambda_1 + 1, \Lambda_2),$	$(\Lambda_1, \Lambda_2 - 1),$	$(\Lambda_1 - 2, \Lambda_2 - 1)$

INTEGRAL REPRESENTATION OF WHITTAKER FUNCTION

§3 Representations of the maximal compact subgroup.

For our later computation, we recall some basic facts about the representation of the maximal compact subgroup K or its complexification $K_{\mathbb{C}}$. Since K is identified with the unitary group of degree 2 $U(2)$, $K_{\mathbb{C}}$ is isomorphic to $GL(2, \mathbb{C})$. Recall a basis of $\mathfrak{u}(2)_{\mathbb{C}}$ given in Section 1:

$$Z = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, H' = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, X = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \bar{X} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

The irreducible finite-dimensional representations of the Lie algebra $\mathfrak{gl}(2, \mathbb{C})$ are parametrized by a set

$$\{\lambda = (\lambda_1, \lambda_2) \in \mathbb{Z}^{\oplus 2} \mid \lambda_1 \geq \lambda_2, \text{ i.e. } \lambda \text{ is dominant}\}.$$

For each dominant weight λ , we set $d = \lambda_1 - \lambda_2 \geq 0$. Then the dimension of the representation space V_{λ} associated to λ is $d + 1$. We can choose a basis $\{v_k \mid 0 \leq k \leq d\}$ in V_{λ} such that the associated representation τ_{λ} is given by

$$\begin{cases} \tau_{\lambda}(Z)v_k = (\lambda_1 + \lambda_2)v_k; \\ \tau_{\lambda}(H')v_k = (2k - d)v_k; \\ \tau_{\lambda}(X)v_k = (k + 1)v_{k+1}; \\ \tau_{\lambda}(\bar{X})v_k = (d + 1 - k)v_{k-1} = \{d - (k - 1)\}v_{k-1}. \end{cases}$$

Since $H'_1 = \frac{1}{2}(Z + H')$ and $H'_2 = \frac{1}{2}(Z - H')$, we have

$$\tau_{\lambda}(H'_1)v_k = (k + \lambda_2)v_k \quad \text{and} \quad \tau_{\lambda}(H'_2)v_k = (-k + \lambda_1)v_k.$$

If it is necessary to refer explicitly to the dominant weight λ , we denote v_k by $v_{\lambda,k}$.

For the adjoint representation of K on \mathfrak{p}_+ , we have an isomorphism $\mathfrak{p}_+ \cong V_{(2,0)}$, and the correspondence of the basis is given by

$$(X_{(0,2)}, X_{(1,1)}, X_{(2,0)}) \longmapsto (v_0, v_1, v_2).$$

Similarly for \mathfrak{p}_- , we have $\mathfrak{p}_- \cong V_{(0,-2)}$, and the identification of the basis is

$$(X_{(-2,0)}, X_{(-1,-1)}, X_{(0,-2)}) = (v_0, -v_1, v_2).$$

Let us consider a tensor product $V_{\lambda} \otimes \mathfrak{p}_+$. Then it has a decomposition into irreducible factors:

$$V_{\lambda} \otimes \mathfrak{p}_+ \cong V_{\lambda} \otimes V_{(2,0)} = V_{(\lambda_1+2, \lambda_2)} \oplus V_{(\lambda_1+1, \lambda_2+1)} \oplus V_{(\lambda_1, \lambda_2+2)}.$$

Let $P^{(2,0)}$, $P^{(1,1)}$, and $P^{(0,2)}$ be the projectors from $V_{\lambda} \otimes \mathfrak{p}_+$ to the factors $V_{(\lambda_1+2, \lambda_2)}$, $V_{(\lambda_1+1, \lambda_2+1)}$, and $V_{(\lambda_1, \lambda_2+2)}$, respectively. We denote $v_{(2,0),k}$ ($k = 0, 1, 2$) by w_k ($k = 0, 1, 2$).

Lemma (3.1). Set $\mu = (\lambda_1 + 2, \lambda_2)$. Then up to scalars, the projector $P^{(2,0)}$ is given by

$$\begin{aligned} \text{(i)} \quad & P^{(2,0)}(v_{\lambda,k} \otimes w_2) = \frac{(k+1) \cdot (k+2)}{2} v_{\mu,k+2}; \\ \text{(ii)} \quad & P^{(2,0)}(v_{\lambda,k} \otimes w_1) = (k+1)(d+1-k) v_{\mu,k+1}; \\ \text{(iii)} \quad & P^{(2,0)}(v_{\lambda,k} \otimes w_0) = \frac{(d+1-k)(d+2-k)}{2} v_{\mu,k}. \end{aligned}$$

Lemma (3.2). Set $\nu = (\lambda_1 + 1, \lambda_2 + 1)$. Then up to scalars, the projector $P^{(1,1)}$ is given by

$$\begin{aligned} \text{(0)} \quad & P^{(1,1)}(v_{\lambda,d} \otimes w_2) = 0 \\ \text{(i)} \quad & P^{(1,1)}(v_{\lambda,k} \otimes w_2) = (k+1) v_{\nu,k+1} \quad (0 \leq k \leq d-1); \\ \text{(ii)} \quad & P^{(1,1)}(v_{\lambda,k} \otimes w_1) = (d-2k) v_{\nu,k} \quad (0 \leq k \leq d); \\ \text{(iii)} \quad & P^{(1,1)}(v_{\lambda,k} \otimes w_0) = -(d+1-k) v_{\nu,k-1} \quad (1 \leq k \leq d). \end{aligned}$$

Lemma (3.3). Set $\pi = (\lambda_1, \lambda_2 + 2)$. Then up to scalars, the projector $P^{(0,2)}$ is given by

$$\begin{aligned} \text{(i)} \quad & P^{(0,2)}(v_{\lambda,k} \otimes w_2) = v_{\pi,k} \quad (0 \leq k \leq d-2); \\ \text{(ii)} \quad & P^{(0,2)}(v_{\lambda,k} \otimes w_1) = -2 \cdot v_{\pi,k-1} \quad (1 \leq k \leq d-1); \\ \text{(iii)} \quad & P^{(0,2)}(v_{\lambda,k} \otimes w_0) = v_{\pi,k-2} \quad (2 \leq k \leq d); \\ \text{(iv)} \quad & P^{(0,2)}(v_d \otimes w_2) = P^{(0,2)}(v_d \otimes w_1) = P^{(0,2)}(v_{d-1} \otimes w_2) = 0. \end{aligned}$$

Proofs of the above lemmas are easy. It is enough to find the highest weight vectors in $V_\lambda \otimes \mathfrak{p}_+$ corresponding to the factors V_μ , V_ν , and V_π , respectively. Other steps of proofs are settled by induction.

INTEGRAL REPRESENTATION OF WHITTAKER FUNCTION

§4 Characters of the unipotent radical.

Put $N = \exp(\mathfrak{n})$. Then N is written as

$$N = \left\{ \left(\begin{array}{cc|cc} 1 & n_0 & & \\ 0 & 1 & & \\ \hline & & 1 & 0 \\ & & -n_0 & 1 \end{array} \right) \cdot \left(\begin{array}{cc|cc} & & n_1 & n_2 \\ & & n_2 & n_3 \\ \hline & & & \\ & & & 1_2 \end{array} \right) \mid n_0, n_1, n_2, n_3 \in \mathbb{R} \right\}.$$

The commutator group $[N, N]$ of N is given by

$$[N, N] = \left\{ \left(\begin{array}{cc|cc} & & n_1 & n_2 \\ & & n_2 & 0 \\ \hline & & & \\ & & & 1_2 \end{array} \right) \mid n_1, n_2 \in \mathbb{R} \right\}.$$

Hence a unitary character η of N is written as

$$\left(\begin{array}{cc|cc} 1 & n_0 & & \\ 0 & 1 & & \\ \hline & & 1 & 0 \\ & & -n_0 & 1 \end{array} \right) \cdot \left(\begin{array}{cc|cc} & & n_1 & n_2 \\ & & n_2 & n_3 \\ \hline & & & \\ & & & 1_2 \end{array} \right) \mapsto \exp\{2\pi i(c_0 n_0 + c_3 n_3)\}$$

for some real numbers $c_0, c_3 \in \mathbb{R}$.

We denote by the same letter η , the derivative of η

$$\eta : \mathfrak{n} \longrightarrow \mathbb{C}.$$

Since $[\mathfrak{n}, \mathfrak{n}] = \mathbb{R}E_{(e_1 - e_2)} \oplus \mathbb{R}E_{(2e_2)}$, η is determined by the purely imaginary numbers

$$\eta_0 = \eta(E_{e_1 - e_2}) \quad \text{and} \quad \eta_3 = \eta(E_{2e_2}).$$

§5 Characterization of the minimal K -type.

Let $\eta : N = \exp(\mathfrak{n}) \rightarrow \mathbb{C}^*$ be a unitary character. Then we denote by $C_\eta^\infty(N \backslash G)$ the space

$$C_\eta^\infty(N \backslash G) = \{\phi : G \rightarrow \mathbb{C}, C^\infty\text{-function} \mid \phi(ng) = \eta(n)\phi(g), (n, g) \in N \times G\}.$$

By the right regular action of G , $C_\eta^\infty(N \backslash G)$ has structures of smooth G -module, and $(\mathfrak{g}_\mathbb{C}, K)$ -module.

For any finite-dimensional K -module (τ, V) , we put

$$\begin{aligned} C_{\eta, \tau}^\infty(N \backslash G/K) \\ = \{F : G \rightarrow V, C^\infty\text{-function} \mid F(n g k^{-1}) = \eta(n)\tau(k)F(g), (n, g, k) \in N \times G \times K\}. \end{aligned}$$

Let (π_Λ, E_Λ) be the representation of discrete series with Harish-Chandra parameter Λ , and let $(\pi_\Lambda^*, E_\Lambda^*)$ be its contragradient representation.

Assume that there exists a continuous homomorphism $W : (\pi_\Lambda^*, E_\Lambda^*) \rightarrow C_\eta^\infty(N \backslash G)$ of smooth G -modules. Then the restriction of W to the minimal K -type τ_λ^* of π_Λ^* gives an element $F_W \in C_{\eta, \tau_\lambda}^\infty(N \backslash G/K)$ such that

$$W(v^*) = \langle v^*, F_W(\cdot) \rangle \quad \text{for all } v^* \in V_\lambda^*.$$

There is a characterization of the minimal K -type function F by means of a differential operator acting on $C_{\eta, \tau_\lambda}^\infty(N \backslash G/K)$.

Let $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ be a Cartan decomposition of \mathfrak{g} , and $Ad = Ad_{\mathfrak{p}_\mathbb{C}}$ the adjoint representation of K on $\mathfrak{p}_\mathbb{C}$. Then we have a canonical covariant differential operator $\nabla_{\lambda, \eta}$ from $C_{\eta, \tau_\lambda}^\infty(N \backslash G/K)$ to $C_{\eta, \tau_\lambda \otimes Ad}^\infty(N \backslash G/K)$:

$$\nabla_{\eta, \lambda} F = \sum_i L_{X_i} F(\cdot) \otimes X_i, \quad F \in C_{\eta, \tau_\lambda}^\infty(N \backslash G/K),$$

where $(X_i)_i$ is any fixed orthonormal basis of \mathfrak{p} with respect to the Killing form of \mathfrak{g} , and $L_{X_i} F(g) = \left(\frac{d}{dt} F(g \cdot \exp(tX_i)) \right) \Big|_{t=0}$ ($g \in G$).

Let $(\tau_\lambda^-, V_\lambda^-)$ be the sum of irreducible K -submodules of $V_\lambda \otimes \mathfrak{p}_\mathbb{C}$ with highest weights of the form $\lambda - \beta$, β being a non-compact root in Σ^+ . Denote by P_λ a surjective K -homomorphism from $V_\lambda \otimes \mathfrak{p}_\mathbb{C}$ to V_λ^- . We define $\mathcal{D}_{\eta, \lambda}$ as the composite of $\nabla_{\eta, \lambda}$ with P_λ :

$$\begin{aligned} \mathcal{D}_{\eta, \lambda} : C_{\eta, \tau_\lambda}^\infty(N \backslash G/K) &\longrightarrow C_{\eta, \tau_\lambda^-}^\infty(N \backslash G/K), \quad \mathcal{D}_{\eta, \lambda} F = P_\lambda(\nabla_{\eta, \lambda} F(\cdot)) \\ &\quad (F \in C_{\eta, \tau_\lambda}^\infty(N \backslash G/K)). \end{aligned}$$

We have the following

Proposition (5.1). (Proposition (2.1) of Yamashita [Y-I])

Let π_Λ be a representation of discrete series with Harish-Chandra parameter Λ of $\mathrm{Sp}(2, \mathbb{R})$. Set $\lambda = \Lambda - \rho_c + \rho_n$. Then the linear map

$$W \in \mathrm{Hom}_{(\mathfrak{g}_\mathbb{C}, K)}(\pi_\Lambda^*, C_\eta^\infty(N \backslash G)) \longrightarrow F_W \in \mathrm{Ker}(\mathcal{D}_{\eta, \lambda})$$

INTEGRAL REPRESENTATION OF WHITTAKER FUNCTION

is injective.

Let $\text{Hom}_G^\infty(\pi_\Lambda^*, C_\eta^\infty(N \setminus G))$ be the continuous homomorphisms of smooth G -modules, then we have a canonical injection.

$$\text{Hom}_G^\infty(\pi_\Lambda^*, C_\eta^\infty(N \setminus G)) \longrightarrow \text{Hom}_{(\mathfrak{g}_\mathbb{C}, K)}(\pi_\Lambda^*, C_\eta^\infty(N \setminus G)).$$

By the results of Kostant [10], we have

$$\dim_{\mathbb{C}} \text{Hom}_G^\infty(\pi_\Lambda^*, C_\eta^\infty(N \setminus G)) + \dim_{\mathbb{C}} \text{Hom}_G^\infty(\pi_\Lambda, C_\eta^\infty(N \setminus G)) \leq 1,$$

and

$$\dim_{\mathbb{C}} \text{Hom}_{(\mathfrak{g}_\mathbb{C}, K)}(\pi_\Lambda^*, C_\eta^\infty(N \setminus G)) + \dim_{\mathbb{C}} \text{Hom}_{(\mathfrak{g}_\mathbb{C}, K)}(\pi_\Lambda, C_\eta^\infty(N \setminus G)) = 0 \quad \text{or} \quad |W|.$$

Here $|W|$ is the order of the Weyl group of $\text{Sp}(2, \mathbb{R})$, hence 8.

Since holomorphic discrete series and antiholomorphic discrete series are not large in the sense of Vogan [21], if $\pi_\Lambda \in \Xi_I^\cup \Xi_{VI}$, we have

$$\dim_{\mathbb{C}} \text{Hom}_G^\infty(\pi_\Lambda^*, C_\eta^\infty(N \setminus G)) = \dim_{\mathbb{C}} \text{Hom}_{(\mathfrak{g}_\mathbb{C}, K)}(\pi_\Lambda^*, C_\eta^\infty(N \setminus G)) = 0.$$

In subsequent sections, we show that if $\Lambda \in \Xi_{II} \cup \Xi_{III}$,

$$\dim \text{Hom}_{(\mathfrak{g}_\mathbb{C}, K)}(\pi_\Lambda^*, C_\eta^\infty(N \setminus G)) = \dim_{\mathbb{C}} \text{Ker}(\mathcal{D}_{\eta, \lambda}) = \frac{1}{2}|W| = 4,$$

and accordingly

$$\dim_{\mathbb{C}} \text{Hom}_G^\infty(\pi_\Lambda^*, C_\eta^\infty(N \setminus G)) + \dim_{\mathbb{C}} \text{Hom}_G^\infty(\pi_\Lambda, C_\eta^\infty(N \setminus G)) = 1$$

by a theorem of Kostant [10].

§6 Radial part of differential operators.

Put $A = \exp(\mathfrak{a})$, i.e.

$$A = \left\{ \begin{pmatrix} a_1 & & & \\ & a_2 & & \\ & & a_1^{-1} & \\ & & & a_2^{-1} \end{pmatrix} \mid a_1, a_2 \in \mathbb{R}, a_1 > 0, a_2 > 0 \right\}.$$

Then we have the Iwasawa decomposition of $Sp(2; \mathbb{R}) : G = NAK$. The value of $F \in C_{\eta, \tau_\lambda}^\infty(N \backslash G/K)$ is determined by its restriction $\phi = F|_A$ to A .

We compute the radial parts $R(\nabla_{\eta, \lambda})$ and $R(\mathcal{D}_{\eta, \lambda})$ of $\nabla_{\eta, \lambda}$ and $\mathcal{D}_{\eta, \lambda}$, respectively.

As an orthogonal basis of \mathfrak{p} , we take

$$C\|\beta\|(X_\beta + X_{-\beta}), \quad \frac{C\|\beta\|}{\sqrt{-1}}(X_\beta - X_{-\beta}) \quad (\beta \in \Sigma^+)$$

with some $C > 0$ depending on the Killing form. Then

$$2\nabla_{\eta, \lambda} F = C \sum_{\beta \in \Sigma_n^+} \|\beta\|^2 L_{X_{-\beta}} F \otimes X_\beta + C \sum_{\beta \in \Sigma_n^+} \|\beta\|^2 L_{X_\beta} F \otimes X_{-\beta}.$$

We write

$$\begin{cases} \nabla_{\eta, \lambda}^+ F = \frac{1}{4} \Sigma \|\beta\|^2 \cdot L_{X_{-\beta}} F \otimes X_\beta; \\ \nabla_{\eta, \lambda}^- F = \frac{1}{4} \Sigma \|\beta\|^2 \cdot L_{X_\beta} F \otimes X_{-\beta}. \end{cases}$$

In order to write $R(\nabla_{\eta, \lambda}^\pm)$, it is better to introduce some “macro” symbols. We set $\partial_i = L_{H_i}$ restricted to A . ($i = 1, 2$), and define linear differential operators \mathcal{L}_i^\pm and S^\pm on $C^\infty(A, V_\lambda)$ by

$$\begin{cases} \mathcal{L}_i^\pm \phi = (\partial_i \pm 2\sqrt{-1} a_i^2 \eta(E_{2e_i}))\phi \quad (i = 1, 2); \\ S^\pm \phi = \{a_1 a_2^{-1} \eta(E_{e_1 - e_2}) \pm \sqrt{-1} a_1 a_2 \eta(E_{e_1 + e_2})\} \phi. \end{cases}$$

Proposition (6.1). *The operators $R(\nabla_{\eta, \lambda}^\pm) = C^\infty(A, V_\lambda) \rightarrow C^\infty(A, V_\lambda \otimes \mathfrak{p}_\pm)$ are expressed as*

$$\begin{aligned} (i) \quad R(\nabla_{\eta, \lambda}^+) \phi &= (\mathcal{L}_1^- + \tau_\lambda \otimes \text{Ad}_{\mathfrak{p}_+}(H'_1) - 4)(\phi \otimes X_{(2,0)}) \\ &\quad + (S^- + \tau_\lambda \otimes \text{Ad}_{\mathfrak{p}_+}(X))(\phi \otimes X_{(1,1)}) \\ &\quad + (\mathcal{L}_2^- + \tau_\lambda \otimes \text{Ad}_{\mathfrak{p}_+}(H'_2) - 2)(\phi \otimes X_{(0,2)}) \\ (ii) \quad R(\nabla_{\eta, \lambda}^-) \phi &= (\mathcal{L}_1^+ - \tau_\lambda \otimes \text{Ad}_{\mathfrak{p}_-}(H'_1) - 4)(\phi \otimes X_{(-2,0)}) \\ &\quad + (S^+ - \tau_\lambda \otimes \text{Ad}_{\mathfrak{p}_-}(\bar{X}))(\phi \otimes X_{(-1,-1)}) \\ &\quad + (\mathcal{L}_2^+ - \tau_\lambda \otimes \text{Ad}_{\mathfrak{p}_-}(H'_2) - 2)(\phi \otimes X_{(0,-2)}). \end{aligned}$$

Proof) In order to prove (i), we have to note that

$$\begin{aligned} (L_{X_{-(2,0)}} F)|_A \otimes X_{(2,0)} &= \{-H'_1 + H_1 - 2\sqrt{-1} E_{2e_1}\} F|_A \otimes X_{(2,0)} \\ &= \{\mathcal{L}_1^- \phi + (\tau_\lambda(H'_1) \cdot F)|_A\} \otimes X_{(2,0)}, \end{aligned}$$

and

INTEGRAL REPRESENTATION OF WHITTAKER FUNCTION

$$\begin{aligned}
(\tau_\lambda(H'_1) \cdot F)|_A \otimes X_{(2,0)} &= \tau_\lambda \otimes Ad_{\mathfrak{p}_+}(H'_1)(\phi \otimes X_{(2,0)}) - \phi \otimes [H'_1, X_{(2,0)}] \\
&= \tau_\lambda \otimes Ad_{\mathfrak{p}_+}(H'_1)(\phi \otimes X_{(2,0)}) - 2(\phi \otimes X_{(2,0)}).
\end{aligned}$$

The case of (ii) is similar.

(q.e.d.)

For a non-compact positive root $\beta = (\beta_1, \beta_2)$ in Σ^+ , let P^β be the projector from $V_\lambda \otimes \mathfrak{p}_+$ to $V_{\lambda+\beta}$, and $P^{-\beta}$ the projector from $V_\lambda \otimes \mathfrak{p}_-$ to $V_{\lambda-\beta}$.

Then, similarly as Lemma (5.2) of Yamashita [I], we can show the following.

Lemma (6.2). *Let λ be the minimal K -type of a discrete series representation π_Λ with Harish-Chandra parameter Λ .*

(i) When $\Lambda \in \Xi_{II}$, $R(\mathcal{D}_{\eta,\lambda})\phi = 0$ if and only if

$$\begin{cases} P^{(0,2)}(R(\nabla_{\eta,\lambda}^+)\phi) = 0; \\ P^{(-1,-1)}(R(\nabla_{\eta,\lambda}^-)\phi) = 0; \\ P^{(-2,0)}(R(\nabla_{\eta,\lambda}^-)\phi) = 0. \end{cases}$$

(ii) When $\Lambda \in \Xi_{III}$, $R(\mathcal{D}_{\eta,\lambda})\phi = 0$ if and only if

$$\begin{cases} P^{(1,1)}(R(\nabla_{\eta,\lambda}^+)\phi) = 0; \\ P^{(0,2)}(R(\nabla_{\eta,\lambda}^+)\phi) = 0; \\ P^{(-2,0)}(R(\nabla_{\eta,\lambda}^-)\phi) = 0. \end{cases}$$

§7 Difference-differential equations.

In this section, we write the system of differential equations in the last lemma of the previous section explicitly in terms of components of ϕ .

Let $\lambda = (\lambda_1, \lambda_2)$ be the minimal K -type of the discrete series representation π_λ . Then in V_λ , we choose a basis $\{v_k \mid 0 \leq k \leq d\}$ defined in Section 3. Here $d = \lambda_1 - \lambda_2$. Then $\phi : A \rightarrow V_\lambda$ is written as

$$\phi(a) = \sum_{k=0}^d c_k(a) v_k$$

with coefficients $c_k(a) : A \rightarrow \mathbb{C}$.

Lemma (7.1).

(i) The condition $P^{(1,1)}(R(\nabla_{\eta,\lambda}^+) \phi) = 0$ is equivalent to the system:

$$\begin{aligned} (C_2^+)_k : & k(\mathcal{L}_1^- + \lambda_2 + d - k - 1)c_{k-1}(a) + (d - 2k)\mathcal{S}^- c_k(a) \\ & + (k - d)(\mathcal{L}_2^- + \lambda_1 - k - 1)c_{k+1}(a) = 0 \quad (0 \leq k \leq d). \end{aligned}$$

(ii) The condition $P^{(-1,-1)}(R(\nabla_{\eta,\lambda}^+) \phi) = 0$ is equivalent to the system:

$$\begin{aligned} (C_2^-)_k : & (k - d)(\mathcal{L}_1^+ - \lambda_2 + k - d - 1)c_{k+1}(a) + (2k - d)\mathcal{S}^+ c_k(a) \\ & + k(\mathcal{L}_2^+ - \lambda_1 + k - 1)c_{k-1}(a) = 0 \quad (0 \leq k \leq d). \end{aligned}$$

(iii) The condition $P^{(0,2)}(R(\nabla_{\eta,\lambda}^+) \phi) = 0$ is equivalent to the system:

$$\begin{aligned} (C_3^+)_k : & (\mathcal{L}_1^- + \lambda_2 - k - 2)c_k(a) - 2\mathcal{S}^- c_{k+1}(a) \\ & + (\mathcal{L}_2^- + \lambda_1 - k - 2)c_{k+2}(a) = 0 \quad (0 \leq k \leq d - 2). \end{aligned}$$

(iv) The condition $P^{(0,-2)}(R(\nabla_{\eta,\lambda}^-) \phi) = 0$ is equivalent to the system:

$$\begin{aligned} (C_3^-)_k : & (\mathcal{L}_1^+ - \lambda_2 - 2d + k)c_{k+2}(a) + 2\mathcal{S}^+ c_{k+1}(a) \\ & + (\mathcal{L}_2^+ - \lambda_1 + k)c_k(a) = 0 \quad (0 \leq k \leq d - 2). \end{aligned}$$

Here we interpret, in the above formulae, $c_{-1}(a) = c_{d+1}(a) = 0$.

Proof) The proof is a direct computation and easy. We omit it.

Since η is trivial on the commutator group $[N, N]$, we have

$$\begin{cases} \mathcal{L}_1^+ = \mathcal{L}_1^- = \partial_1 = a_1 \frac{\partial}{\partial a_1} \\ \mathcal{S}^+ = \mathcal{S}^- = \frac{a_1}{a_2} \eta(E_{(1,-1)}). \end{cases}$$

Thus we have the following

Proposition (7.2). Under the same assumption as in Lemma (6.2) (i), $\phi(a) = \sum_{k=0}^d c_k(a) v_k$ satisfies the following system of partial differential equations:

$$\begin{aligned} (C_3^+)_k : & (\mathcal{L}_1 + \lambda_2 - k - 2)c_k(a) - 2\mathcal{S} \cdot c_{k+1}(a) \\ & + (\mathcal{L}_2^- + \lambda_1 - k - 2)c_{k+2}(a) = 0 \quad (0 \leq k \leq d - 2) \\ (C_3^-)_k : & (\mathcal{L}_1 - \lambda_2 - 2d + k)c_{k+2}(a) + 2\mathcal{S} \cdot c_{k+1}(a) \\ & + (\mathcal{L}_2^+ - \lambda_1 + k)c_k(a) = 0 \quad (0 \leq k \leq d - 2) \\ (C_2^-)_{k+1} : & (k + 1 - d)(\mathcal{L}_1 - \lambda_2 - d + k)c_{k+2}(a) + (2k + 2 - d)\mathcal{S} \cdot c_{k+1}(a) \\ & + (k + 1)(\mathcal{L}_2^+ - \lambda_1 + k)c_k(a) = 0 \quad (-1 \leq k \leq d - 1) \end{aligned}$$

(q.e.d.)

INTEGRAL REPRESENTATION OF WHITTAKER FUNCTION

§8 Reduction of the system of partial differential equations.

In this section we reduce the system of partial differential equations of the previous proposition to a simpler holonomic system, when η is generic.

In the first place, we see that the functions $c_k(a)$ is determined by the coefficient of the highest weight vector $c_d(a)$.

In fact, when $k = 0$, or $k = d$

$$\begin{aligned} (C_2^-)_0 : & \quad (\mathcal{L}_1 - \lambda_2 - d - 1)c_1(a) + \mathcal{S}c_0(a) = 0; \\ (C_2^-)_d : & \quad \mathcal{S}c_d(a) + (\mathcal{L}_2^+ - \lambda_1 + d - 1)c_{d-1}(a) = 0. \end{aligned}$$

Moreover for $1 \leq k \leq d-1$, the computation of $(k+1)(C_3^-)_k - (C_2^-)_{k+1}$ yields

$$(\mathcal{L}_1 - \lambda_2 - d - 1)c_{k+2}(a) + \mathcal{S}c_{k+1}(a) = 0.$$

Noting $\lambda_2 + d = \lambda_1$ together with $(C_2^-)_0$, we have

$$(E)_k : \quad (\mathcal{L}_1 - \lambda_1 - 1)c_{k+2}(a) + \mathcal{S}c_{k+1}(a) = 0 \quad (-1 \leq k \leq d-1).$$

Hence $c_0(a), c_1(a), \dots, c_{d-1}(a)$ are determined downward recursively by $c_d(a)$.

The system of the equations (C_2^-) are now replaced by the above $(E)_k$ and

$$(C_2^-)_{d-1} : \quad \mathcal{S}c_d(a) + (\mathcal{L}_2^+ - \lambda_1 + d - 1)c_{d-1}(a) = 0.$$

Thus the system of the equations of Proposition (7.2) in Section 7 is equivalent to a system of equations:

$$\begin{aligned} (F-1) : & \quad (\mathcal{L}_1 - \lambda_1 - 1)c_d(a) + \mathcal{S}c_{d-1}(a) = 0; \\ (F-2) : & \quad (\mathcal{L}_1 - \lambda_1 - 1)c_{d-1}(a) + \mathcal{S}c_{d-2}(a) = 0; \\ (F-3) : & \quad \mathcal{S}c_d(a) + (\mathcal{L}_2^+ - \lambda_1 + d - 1)c_{d-1}(a) = 0; \\ (F-4) : & \quad (\mathcal{L}_1 + \lambda_2 - d)c_{d-2}(a) - 2\mathcal{S}c_{d-1}(a) + (\mathcal{L}_2^- + \lambda_1 - d)c_d(a) = 0. \end{aligned}$$

In order to make the above equations simpler, we replace unknown functions $c_k(a)$ by $h_k(a)$ defined by relations

$$c_k(a) = a_1^{\lambda_1+1-d} a_2^{\lambda_1} \left(\frac{a_1}{a_2} \right)^k e^{-i\eta(E_{(0,2)})a_2^2} h_k(a)$$

Now we introduce Euler operators ∂_i ($i = 1, 2$) by $\partial_i = a_i \frac{\partial}{\partial a_i}$ for each $i = 1, 2$. Then the system of equations $(F-1) \sim (F-4)$ is replaced by

$$\begin{aligned} (G-1) : & \quad \partial_1 h_d(a) + \eta(E_{e_1-e_2})h_{d-1}(a) = 0; \\ (G-2) : & \quad (\partial_1 - 1)h_{d-1}(a) + \eta(E_{e_1-e_2})h_{d-2}(a) = 0; \\ (G-3) : & \quad \mathcal{S} \left(\frac{a_1}{a_2} \right) h_d(a) + \partial_2 h_{d-1}(a) = 0; \\ (G-4) : & \quad (\partial_1 + 2\lambda_2 - 1)h_{d-2}(a) - 2\mathcal{S} \left(\frac{a_1}{a_2} \right) h_{d-1}(a) \\ & \quad + \left(\frac{a_1}{a_2} \right)^2 (\partial_2 + 2\lambda_1 - 2d - 2\mathcal{S}')h_d(a) = 0. \end{aligned}$$

TAKAYUKI ODA

Here $S' = \frac{1}{2}(\mathcal{L}_2^+ - \mathcal{L}_2^-) = 2\sqrt{-1}\eta(E_{2e_2})a_2^2$.

(G-1) and (G-3) are equivalent to a single equation:

$$(H-1): (\partial_1 \partial_2 - S^2)h_d(a) = 0.$$

(G-1), (G-2), and (G-4) is equivalent to a single equation:

$$(*) : (\partial_1 + 2\lambda_2 - 1)(\partial_1 - 1)\partial_1 \left\{ \left(\frac{a_1}{a_2} \right)^2 h_d(a) \right\} + 2 \left(\frac{a_1}{a_2} \right)^2 \partial_1 h_d(a) \\ + \left(\frac{a_1}{a_2} \right)^2 (\partial_2 + 2\lambda_2 - 2S')h_d(a) = 0.$$

Here we used the assumption that η is generic, i.e.

$$\eta(E_{e_1-e_2}) = \eta_0 \neq 0, \quad \text{and} \quad \eta(E_{2e_2}) = \eta_3 \neq 0.$$

Apply the operator ∂_2 to the above equation (*), and use (H-1) to replace $\partial_1 \partial_2 h_d(a)$ by $S^2 h_d(a)$. Then we have

$$(H-2): \{\partial_1^2 + 2\partial_1 \partial_2 + \partial_2^2 + (2\lambda_2 - 2)(\partial_1 + \partial_2) + (-2\lambda_2 + 1) - 2S' \partial_2\}h_d = 0$$

At last we have the following

Lemma (8.1). *The system of equations of Proposition (7.2) is equivalent to*

$$(H-1): (\partial_1 \partial_2 - S^2)h_d(a) = 0$$

and

$$(H-2): \{(\partial_1 + \partial_2)^2 + (2\lambda_2 - 2)(\partial_1 + \partial_2) + (-2\lambda_2 + 1) - 2S' \partial_2\}h_d(a) = 0$$

We can easily check that the system (H-1), (H-2) is a holonomic system of rank 4 defined over $(\mathbb{R}_{>0})^2 = \{(a_1, a_2) \in \mathbb{R}^2 \mid a_1, a_2 > 0\}$. Hence $\dim_{\mathbb{C}} \text{Ker}(D_{\eta, \lambda}) = 4$. The contragradient representation π_{Λ}^* of π_{Λ} ($\Lambda \in \Xi_{\text{II}}$) is written as $\pi_{\Lambda}^* = \pi_{\Lambda'}$ with some $\Lambda' \in \Xi_{\text{III}}$. Using the difference-differential equations (C_2^+) , (C_3^+) and (C_3^-) , we can similarly show that $\dim_{\mathbb{C}} \text{Ker}(D_{\eta, \lambda'}) = 4$ for the minimal K -type λ' of $\pi_{\Lambda'}$.

Since Kostant's result implies (cf. §5)

$$8 = \dim \text{Hom}_{(\mathfrak{g}_{\mathbb{C}}, K)}(\pi_{\Lambda}^*, c_{\eta}^{\infty}(N \backslash G)) + \dim \text{Hom}_{(\mathfrak{g}_{\mathbb{C}}, K)}(\pi_{\Lambda}, c_{\eta}^{\infty}(N \backslash G)) \\ \leq \dim_{\mathbb{C}} \text{Ker}(D_{\eta, \lambda}) + \dim_{\mathbb{C}} \text{Ker}(D_{\eta, \lambda'}) = 8,$$

we have the following

Proposition. *Assume that η is generic, i.e.*

$$\eta_0 = \eta(E_{e_1-e_2}) \neq 0 \quad \text{and} \quad \eta_3 = \eta(E_{2e_2}) \neq 0.$$

Then for a discrete series representation π_{Λ} ($\Lambda \in \Xi_{\text{II}} \cup \Xi_{\text{III}}$),

$$\dim_{\mathbb{C}} \text{Hom}_{(\mathfrak{g}, K)}(\pi_{\Lambda}, c_{\eta}^{\infty}(N \backslash G)) = 4.$$

INTEGRAL REPRESENTATION OF WHITTAKER FUNCTION

When $\Lambda \in \Xi_{II} \cup \Xi_{III}$, we have

$$\dim_{\mathbb{C}} \text{Hom}_G^\infty(\pi_\Lambda^*, c_\eta^\infty(N \setminus G)) + \dim_{\mathbb{C}} \text{Hom}_G^\infty(\pi_\Lambda, c_\eta^\infty(N \setminus G)) = 1.$$

Hence two cases occur:

$$(A) \quad \text{Hom}_G^\infty(\pi_\Lambda^*, c_\eta^\infty(N \setminus G)) \cong \mathbb{C}, \quad \text{and} \quad \text{Hom}_G^\infty(\pi_\Lambda, c_\eta^\infty(N \setminus G)) = \{0\}$$

or

$$(B) \quad \text{Hom}_G^\infty(\pi_\Lambda^*, c_\eta^\infty(N \setminus G)) = \{0\}, \quad \text{and} \quad \text{Hom}_G^\infty(\pi_\Lambda, c_\eta^\infty(N \setminus G)) \cong \mathbb{C}.$$

In the next section, we see that this dichotomy is controlled by the parity of the imaginary part of the purely imaginary number $\eta_3 = \eta(E_{2e_2}) \neq 0$. And when $\text{Hom}_G^\infty(\pi_\Lambda^*, c_\eta^\infty(N \setminus G))$ is non-zero and generated by W , we have an explicit integral formula for the image $F_W \in \text{Ker}(D_{\eta, \lambda}) \subset c_{\eta, \tau_\lambda}^\infty(N \setminus G/K)$ of the intertwining operator W .

§9. Integral formula for the Whittaker function..

Let us recall the confluent hypergeometric equation given by Whittaker ([], Chap.16):

$$\frac{d^2}{dz^2} W + \left\{ -\frac{1}{4} + \frac{k}{z} + \frac{\frac{1}{4} - m^2}{z^2} \right\} W = 0.$$

When $\text{Re}(k - \frac{1}{2} - m) \leq 0$, for $z \notin (-\infty, 0)$, a unique solution, which rapidly decreases if $z \rightarrow +\infty$, is given by

$$W_{k,m}(z) = \frac{e^{-\frac{1}{2}z} \cdot z^k}{\Gamma(\frac{1}{2} - k + m)} \int_0^\infty t^{-k-\frac{1}{2}+m} \left(1 + \frac{t}{z}\right)^{k-\frac{1}{2}+m} \cdot e^{-t} dt.$$

The following is the main result of this paper.

Theorem (9.1). Assume that $\eta : N \rightarrow \mathbb{C}^*$ is generic, i.e. $\eta_0 = \eta(E_{e_1-e_2}) \neq 0$ and $\eta_3 = \eta(E_{2e_2}) \neq 0$.

(i) For $\Lambda \in \Xi_{II}$,

$$\begin{cases} \text{Hom}_G^\infty(\pi_\Lambda^*, c_\eta^\infty(N \setminus G)) \cong \mathbb{C}, & \text{if } \text{Im}(\eta_3) < 0; \\ \text{Hom}_G^\infty(\pi_\Lambda^*, c_\eta^\infty(N \setminus G)) = \{0\}, & \text{if } \text{Im}(\eta_3) > 0. \end{cases}$$

(ii) Assume that $\Lambda \in \Xi_{II}$ and $\text{Im}(\eta_3) < 0$, and let W be a unique intertwining operator in $\text{Hom}_G^\infty(\pi_\Lambda^*, c_\eta^\infty(N \setminus G))$ up to scalar multiple. Then the function $h_d(a_1, a_2)$ associated to $\phi(a) = F_{W|A}(a) = \sum_{i=0}^d c_i(a) v_i$ ($F_W \in \text{Ker}(\mathcal{D}_{\eta, \tau_\lambda})$) has an integral representation

$$h_d(a_1, a_2) = \int_0^\infty t^{\lambda_2 - \frac{3}{2}} W_{0, -\lambda_2}(t) \exp\left(-\frac{t^2}{32\sqrt{-1} \cdot \eta_3 a_2^2} + \frac{8\sqrt{-1}\eta_0^2 \eta_3 a_1^2}{t^2}\right) \frac{dt}{t}$$

Proof. It is easy to check that the integral represents a solution of the differential equations $(H - 1)$ and $(H - 2)$, by derivation of integrand and partial integration.

Replace t by $a_1 t$ in the above integral expression of $h_d(a_1, a_2)$, then

$$h_d(a_1, a_2) = \int_0^\infty \left(\frac{a_1}{a_2} \cdot a_2 \cdot t \right)^{\lambda_2 - \frac{3}{2}} W_{0, -\lambda_2} \left(\frac{a_1}{a_2} \cdot a_2 \cdot t \right) \\ \times \exp \left\{ -\frac{1}{32\sqrt{-1}\eta_3} \left(\frac{a_1}{a_2} \right)^2 \cdot t^2 + 8\sqrt{-1}\eta_0^2\eta_3 \cdot t^{-2} \right\} \frac{dt}{t}.$$

If $\text{Im}(\eta_3) < 0$, $-\frac{1}{32\sqrt{-1}\eta_3} < 0$ and $8\sqrt{-1}\eta_0^2\eta_3 < 0$. Also since $\Lambda \in \Xi_{II}$, λ_2 is a negative integer. Hence the integrand is rapidly decreasing when $t \rightarrow +\infty$, and when $t \rightarrow 0$. Therefore the above integral converges, and as a function in (a_1, a_2) , it is rapidly decreasing when $\frac{a_1}{a_2} \rightarrow \infty$ and $a_2 \rightarrow \infty$. Put

$$c_d(a) = a_1^{\lambda_1+1-d} a_2^{\lambda_1} \cdot \left(\frac{a_1}{a_2} \right)^d \cdot e^{-i\eta_3 \cdot a_2^2} \cdot h_d(a),$$

and $c_k(a)$ for $0 \leq k \leq d-1$ by the recurrence relation $(E)_k$ of §8.

Then $c_k(a)$ ($0 \leq k \leq d$) are also rapidly decreasing functions in $(a_1/a_2, a_2)$. Write $\phi(a) = \sum_{k=0}^d c_k(a) v_k \in C^\infty(A, V_\lambda)$. Then for any vector v^* of the dual space V_λ^* , $(\phi(a), v^*)$ is also a rapidly decreasing function. *A fortiori*, $\phi(a)$, i.e. $F(g) = \eta(n)\tau_\lambda(k)^{-1}\phi(a)$ is slowly increasing in $g = nak \in G$. This F defines an element W in $\text{Hom}_{\mathfrak{g}, K}(\pi_\Lambda^*, C^\infty(N \setminus G))$ by Schmidt's characterization ([]).

Now Wallach's version of multiplicity one (cf. [W], §8) implies that the operators W in $\text{Hom}_{\mathfrak{g}, K}(\pi_\Lambda^*, C^\infty(N \setminus G))$ such that $W(v)$ are slowly increasing on G for any $v \in \pi_\Lambda$, forms a linear subspace of dimension at most one.

Hence $\text{Hom}_G^\infty(\pi_\Lambda^*, C^\infty(N \setminus G)) \neq \{0\}$, if $\text{Im}(\eta_3) < 0$. If $\text{Im}(\eta_3) > 0$, by a similar argument, we can show that $\text{Hom}_G^\infty(\pi_\Lambda, c_\eta^\infty(N \setminus G)) \neq \{0\}$. Since

$$\dim_{\mathbb{C}} \text{Hom}_G^\infty(\pi_\Lambda^*, C^\infty(N \setminus G)) + \dim_{\mathbb{C}} \text{Hom}_G^\infty(\pi_\Lambda, c_\eta^\infty(N \setminus G)) = 1$$

if η is generic, this proves (i). The part (ii) immediately follows from the uniqueness of the Whittaker model.

Remark. For general cases, the condition of (i) is described in terms of wave front set by Matsumoto ([], §).

When $G = SU(2, 2)$, we have a similar integral expression of the Whittaker function of the highest weight vector of the minimal K -type of a discrete series representation. Details are discussed elsewhere about this case.

REFERENCES

- [H-S] Hecht, H. and Schmid, W., *A proof of Blattner's conjecture*, *Inventiones math.* **31** (1975), 129–154.
- [J-L] Jacquet, H. and Langlands, R. P., *Automorphic forms on $GL(2)$* , *Lecture Notes in Math.*, No. 114, Springer.
- [K] Kostant, B., *On Whittaker Vectors and Representation Theory*, *Inventiones Math.* **48** (1978), 101–184.
- [M] Matsumoto, H., *$C^{-\infty}$ -Whittaker vectors corresponding to a principal nilpotent orbit of a real reductive linear group and wave front sets*, Preprint.

INTEGRAL REPRESENTATION OF WHITTAKER FUNCTION

- [S] Schmid, W., *On the realization of the discrete series of a semisimple Lie group*, Rice Univ. Studies **56** (1970), 99–108.
- [V] Vogan, D. Jr., *Gelfand-Kirillov Dimension for Harish-Chandra Modules*, Inventiones Math. **49** (1978), 75–98.
- [W] Wallach, N., *Asymptotic expansions of generalized matrix entries of representations of real reductive groups*, Lie Group Representations I, Lecture Notes in Math. 1024 ((1984)), 287–369.
- [W-W] Whittaker, E.T. and Watson, G.N., *A course of modern analysis*, Cambridge Univ. Press, 4ed., (1965).
- [Y-I] Yamashita, H., *Embedding of discrete series into induced representations of semisimple Lie groups, I*, General theory and the case of for $SU(2, 2)$, Japan J. Math. **16** (1990), 31–95.
- [Y-II] Yamashita, H., *Embedding of discrete series into induced representations of semisimple Lie groups, II*, -Generalized Whittaker models for $SU(2, 2)$ -, J. Math. Kyoto Univ. **31-1** (1991), 543–571.

RIMS, KYOTO UNIV., KYOTO, 606, JAPAN